lution, the mucus separates in the form of a flocculent coagulum, which is tough and stringy, and on drying has a brownish colour. When chewed, it soon fills the mouth with mucilage.

Silicated potash rendered the upper stratum of the liquid dark and opaque, but without any precipitate like that which takes place in the mucilage of gum.

2. Tannin. This is readily thrown down from the decoction and tincture by gelatin. The sulphate of iron strikes with it a very black colour.

3. Resin. This also exists plentifully. It communicates to alcohol a reddish colour, and is instantly precipitated from it by water. When obtained pure, it is of a reddish cast, fusible, inflammable and moderately bitter.

I have not detected any extractive, properly so called, in these leaves. When the muriate of tin is added to the decoction, it separates a very copious yellow precipitate. This however is owing to the mucus. If alcohol be first added to the decoction, and the coagulum which it forms withdrawn; the fluid no longer gives a precipitate with muriate of tin, although it readily yields one to gelatin.

Distillation with water affords a mild fluid with little taste or odour.

The Kalmia latifolia, together with some other species of its genus, has long had the reputation, in various parts of the country, of being poisonous to certain domestic animals. Catesby says of it, that "deer feed on its green leaves with impunity; yet when cattle and sheep, by severe winters deprived of better food, feed on the leaves of this plant, a great many of them die annually."

Kalm, the Swedish traveller, who gave name to this genus, says of Kalmia latifolia, "The leaves are poison to some animals, and food for others; experience has taught the people, that when sheep eat of these leaves, they either die immediately, or fall very sick, and recover with great difficulty. The young and more tender sheep are killed by a small portion, but the older ones can bear a stronger dose. Yet this food will likewise prove mortal to them, if they take too much of it. The same noxious effect it shews in regard to calves which eat too much of the leaves; they either die, or do not recover easily. I can remember that in the year 1748 some calves ate of the leaves; they fell very sick, swelled, foamed at the mouth and could hardly stand; however, they were cured by giving them gunpowder and other medicines. The sheep are most exposed to be tempted with these leaves in winter, for after having been kept in sta-